

# Snapshots of Electrons in Motion


Recent, groundbreaking experiments using ultrashort laser pulses have permitted the study of the motion of electrons in atoms right after ionization. This leads the way to a better understanding and control of the motion of electrons in atoms and molecules.

Technology tends to become faster, by about three orders of magnitude, every decade or so. An international team of researchers from Germany, the United States of America, and Saudi Arabia were recently able to record one of the fastest processes known: the electron motion inside an atom right after ionization. This brings the experimental study of ultrafast attosecond processes in atomic physics and chemistry within the reach of experimental study.

Attosecond research is nowadays at the forefront of technological and scientific progress. One attosecond is 10<sup>-18</sup> seconds – a billionth of a billionth of a second: a truly tiny time interval, indeed. Many chemical and physical processes, ranging from simple electron dynamics in atoms to complex charge migration in biomolecules, happen at such a timescale. The steady development of attosecond technology, a natural consequence of ever-faster electronics and optics, is now opening the doors to the study of some of the fastest known processes happening in such incredibly short intervals of time.

“The basic principles behind attosecond technology are conceptually similar to stop-motion photography,” explains Eleftherios Goulielmakis, who led the experiment at the Max-Planck-Institute for Quantum Optics in Munich (Germany). The image sensor (or film) constantly captures light reflected by a fast-moving object. A popular way amongst photographers to freeze the motion of such objects is to use a short exposure time to reduce the blur, and a flash to concentrate the light reflected by the object, within a short period of time. In the same way, achieving a high temporal resolution is arguably the most powerful way to experimentally study ultrafast processes.

Many groundbreaking ultrafast tools and techniques have been developed in recent years [1]. In particular, “attosecond laser pulses allow us to freeze motion,” explains Adrian Wirth, who worked with Goulielmakis. “First, we eject one electron from a krypton atom by using a strong laser pulse. Then, after a specific time, we freeze the motion of the remaining electrons right after ionization, by using an attosecond laser pulse. In essence, our attosecond laser pulse thus acts just like a very sophisticated flash.” In order to reconstruct the quantum mechanical wavefunction of the remaining electrons, an image is always taken of a large number of synchronized atoms. Finally, the dynamics of the process is reconstructed by combining a number of




**Figure 1: Stop-motion to see the electrons move.**

The ultra-high temporal resolution of 150 attoseconds enables the reconstruction of the evolution of the electron distribution of the krypton atoms right after ionization. Notice that this evolution occurs within a few femtoseconds only. Picture: Christian Hackenberger, LMU.

images taken at different stages of the wavefunction evolution. “The most intuitive way to think of the measurement process,” continues Goulielmakis, “is by realizing that the distribution of the electron cloud around the ion influences how much of the attosecond pulse is absorbed. If the cloud is aligned like an antenna, it absorbs more radiation compared to when it is not. A detector behind the atoms measures the energy of the attosecond pulse at each snapshot and this allows us to reconstruct the motion.”

“This is an important experiment because, for the first time, researchers have been able to resolve the motion of electrons within an atom,” says Raphael Levine [2] from the Hebrew University of Jerusalem (Israel) and the University



**Figure 2: The eye of the camera.** The look and working principle of an ultrafast apparatus differ radically from those of a conventional camera: krypton atoms are placed inside a vacuum chamber where they are ionized with a first laser pulse and subsequently probed by a second, ultra-short laser pulse. Picture: Eleftherios Goulielmakis, MPQ.

of California in Los Angeles (USA). “From a theorist’s point of view,” he continues, “I am, of course, happy to see that their experiment shows our general understanding is accurately describing the physical behavior. Nonetheless, it is important to realize that this research shows we no longer have to stick to Max Born’s 85-year-old paradigm treating electron motion as instantaneous. Now, it is actually possible to see how the electrons move!”

“This work is a perfect demonstration of the enormous potentialities of attosecond science, whose main goal is the real time observation and control of electron motion in atoms, molecules and solids,” says Mauro Nisoli [2] from Politecnico of Milan (Italy). “As a consequence, we can now finally tackle several open theoretical questions experimentally. For example, it now becomes possible to study electron correlation effects or ultrafast multi-electron dynamics. On a more general level, a fundamental open question

is whether the reaction pathways in a chemical reaction can be controlled and steered by controlling the electronic coherent motion on an attosecond timescale. If this turned out to be the case, we would soon see the new field of attochemistry emerge, expanding the current tools available through femto-chemistry.” In fact, femto-chemistry has emerged as a consequence of new femtosecond technology about a decade ago. Today, femto-chemistry has matured into a standard way to influence chemical reactions with light and it is, for example, a powerful tool for laser-based isotope separation.

“Since the 1960s,” Levine recalls, “we have seen rather regular improvements in our capabilities of dealing with ever shorter intervals of time. Back then, we were dealing with microseconds and microelectronics, today with attoseconds and advanced optics — and every three orders of magnitude somebody was awarded with the Nobel Prize.” “Indeed,” Nisoli agrees, “if attosecond technology proves to be useful in accessing the fascinating microcosmos microcosmos of ultra-fast processes, I am also convinced that it will have a tremendous impact on science and technology, and therefore receive its due credit.” Apart from known ultrafast processes, both researchers are convinced that further, unexpected applications are bound to surface, thus expanding the impact of attosecond technology even further. “After all,” Levine concludes, “I have always enjoyed working in this field because it is intellectually very challenging. And experience shows that, sooner or later, interesting science always leads to fascinating new applications.”

[1] A. Niederberger, *Towards Filming Chemical Reactions*, *Opt. Photon. Focus* **2**, 6 (2008). <http://www.opfocus.org/index.php?topic=story&v=2&s=6>

[2] F. Remacle & R. D. Levine, *An electronic time scale in chemistry*, *PNAS* **103**, 6793-6798 (2006).

[3] G. Sansone *et al.*, *Electron localization following attosecond molecular photoionization*, *Nature* **465**, 763-766 (2010).

**Armand Niederberger**  
© 2010 Optics & Photonics Focus

Eleftherios Goulielmakis, Zhi-Heng Loh, Adrian Wirth, Robin Santra, Nina Rohringer, Vladislav S. Yakovlev, Sergey Zharebtsov, Thomas Pfeifer, Abdallah M. Azzeer, Matthias F. Kling, Stephen R. Leone & Ferenc Krausz, **Real-time observation of valence electron motion**, *Nature* (2010) **466**, 739-743.